

The **Equation of Energy** in Cartesian, cylindrical, and spherical coordinates for

Newtonian fluids of constant density, with source term S_e . Source could be electrical energy due to current flow, chemical energy, etc. Two cases are presented: the general case where thermal conductivity may be a function of temperature (vector flux $\tilde{q} = q/\text{area}$ appears in the equations); and the more usual case, where thermal conductivity is constant (on the reverse).

Spring 2020 Faith A. Morrison, Michigan Technological University

Microscopic energy balance, in terms of flux; Gibbs notation

$$\rho \hat{C}_p \left(\frac{\partial T}{\partial t} + \underline{v} \cdot \nabla T \right) = -\nabla \cdot \underline{\tilde{q}} + S_e$$

Microscopic energy balance, in terms of flux; Cartesian coordinates

$$\rho \hat{C}_p \left(\frac{\partial T}{\partial t} + v_x \frac{\partial T}{\partial x} + v_y \frac{\partial T}{\partial y} + v_z \frac{\partial T}{\partial z} \right) = - \left(\frac{\partial \tilde{q}_x}{\partial x} + \frac{\partial \tilde{q}_y}{\partial y} + \frac{\partial \tilde{q}_z}{\partial z} \right) + S_e$$

Microscopic energy balance, in terms of flux; cylindrical coordinates

$$\rho \hat{C}_p \left(\frac{\partial T}{\partial t} + v_r \frac{\partial T}{\partial r} + \frac{v_\theta}{r} \frac{\partial T}{\partial \theta} + v_z \frac{\partial T}{\partial z} \right) = - \left(\frac{1}{r} \frac{\partial(r\tilde{q}_r)}{\partial r} + \frac{1}{r} \frac{\partial \tilde{q}_\theta}{\partial \theta} + \frac{\partial \tilde{q}_z}{\partial z} \right) + S_e$$

Microscopic energy balance, in terms of flux; spherical coordinates

$$\rho \hat{C}_p \left(\frac{\partial T}{\partial t} + v_r \frac{\partial T}{\partial r} + \frac{v_\theta}{r} \frac{\partial T}{\partial \theta} + \frac{v_\phi}{r \sin \theta} \frac{\partial T}{\partial \phi} \right) = - \left(\frac{1}{r^2} \frac{\partial(r^2 \tilde{q}_r)}{\partial r} + \frac{1}{r \sin \theta} \frac{\partial(\tilde{q}_\theta \sin \theta)}{\partial \theta} + \frac{1}{r \sin \theta} \frac{\partial \tilde{q}_\phi}{\partial \phi} \right) + S_e$$

Fourier's law of heat conduction, Gibbs notation: $\underline{\tilde{q}} = q/A = -k\nabla T$

Fourier's law of heat conduction, Cartesian coordinates: (constant thermal conductivity k)

$$\begin{pmatrix} \tilde{q}_x \\ \tilde{q}_y \\ \tilde{q}_z \end{pmatrix}_{xyz} = \begin{pmatrix} q_x/A \\ q_y/A \\ q_z/A \end{pmatrix}_{xyz} = \begin{pmatrix} -k \frac{\partial T}{\partial x} \\ -k \frac{\partial T}{\partial y} \\ -k \frac{\partial T}{\partial z} \end{pmatrix}_{xyz}$$

Fourier's law of heat conduction, cylindrical coordinates: (constant thermal conductivity k)

$$\begin{pmatrix} \tilde{q}_r \\ \tilde{q}_\theta \\ \tilde{q}_z \end{pmatrix}_{r\theta z} = \begin{pmatrix} q_r/A \\ q_\theta/A \\ q_z/A \end{pmatrix}_{r\theta z} = \begin{pmatrix} -k \frac{\partial T}{\partial r} \\ -\frac{k}{r} \frac{\partial T}{\partial \theta} \\ -k \frac{\partial T}{\partial z} \end{pmatrix}_{r\theta z}$$

Fourier's law of heat conduction, spherical coordinates: (constant thermal conductivity k)

$$\begin{pmatrix} \tilde{q}_r \\ \tilde{q}_\theta \\ \tilde{q}_\phi \end{pmatrix}_{r\theta\phi} = \begin{pmatrix} q_r/A \\ q_\theta/A \\ q_\phi/A \end{pmatrix}_{r\theta\phi} = \begin{pmatrix} -k \frac{\partial T}{\partial r} \\ -\frac{k}{r} \frac{\partial T}{\partial \theta} \\ -\frac{k}{r \sin \theta} \frac{\partial T}{\partial \phi} \end{pmatrix}_{r\theta\phi}$$

The **Equation of Energy** for systems with **constant k**

Microscopic energy balance, constant thermal conductivity; Gibbs notation

$$\rho \hat{C}_p \left(\frac{\partial T}{\partial t} + \underline{v} \cdot \nabla T \right) = k \nabla^2 T + S_e$$

Microscopic energy balance, constant thermal conductivity; Cartesian coordinates

$$\rho \hat{C}_p \left(\frac{\partial T}{\partial t} + v_x \frac{\partial T}{\partial x} + v_y \frac{\partial T}{\partial y} + v_z \frac{\partial T}{\partial z} \right) = k \left(\frac{\partial^2 T}{\partial x^2} + \frac{\partial^2 T}{\partial y^2} + \frac{\partial^2 T}{\partial z^2} \right) + S_e$$

Microscopic energy balance, constant thermal conductivity; cylindrical coordinates

$$\rho \hat{C}_p \left(\frac{\partial T}{\partial t} + v_r \frac{\partial T}{\partial r} + \frac{v_\theta}{r} \frac{\partial T}{\partial \theta} + v_z \frac{\partial T}{\partial z} \right) = k \left(\frac{1}{r} \frac{\partial}{\partial r} \left(r \frac{\partial T}{\partial r} \right) + \frac{1}{r^2} \frac{\partial^2 T}{\partial \theta^2} + \frac{\partial^2 T}{\partial z^2} \right) + S_e$$

Microscopic energy balance, constant thermal conductivity; spherical coordinates

$$\begin{aligned} \rho \hat{C}_p & \left(\frac{\partial T}{\partial t} + v_r \frac{\partial T}{\partial r} + \frac{v_\theta}{r} \frac{\partial T}{\partial \theta} + \frac{v_\phi}{r \sin \theta} \frac{\partial T}{\partial \phi} \right) \\ &= k \left(\frac{1}{r^2} \frac{\partial}{\partial r} \left(r^2 \frac{\partial T}{\partial r} \right) + \frac{1}{r^2 \sin \theta} \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial T}{\partial \theta} \right) + \frac{1}{r^2 \sin^2 \theta} \frac{\partial^2 T}{\partial \phi^2} \right) + S_e \end{aligned}$$

Reference: F. A. Morrison, "Web Appendix to *An Introduction to Fluid Mechanics*," Cambridge University Press, New York, 2013 (pages.mtu.edu/~fmorriso/cm310/IFMWebAppendixDMicroEBalanceMorrison.pdf). This worksheet is on the web at pages.mtu.edu/~fmorriso/cm310/energy.pdf.